



CHARACTERIZATION AND MEASUREMENT OF POLLUTION LEVEL OF GROUND WATER IN THE CONTEXT OF ECOLOGICAL EVALUATION OF SOME PLANT

RAJEEV KUMAR TRIPATHI^a AND VIVEK KUMAR AGRAHARI^{b1}

^{ab}P.G. Department of Botany, R.H.S. P.G. College, Singramau, Jaunpur, Uttar Pradesh, India

ABSTRACT

Water is foremost and legitimate friend of living beings without water life cannot be imagine. The water quality for agriculture, domestic, industrial and commercial purpose. The properties of water such as physical as well as chemical play important role. The deciding the water quality the parameter shows major external effects. The positional and periodic estimation at different site of Karakhion Industrial area of Varanasi district has been done. The physical parameters are compared and there monitoring suggested where whenever alarming me.

KEYWORDS: Evaluation, Foremost, Legitimate, Monitoring, Effluents

The accumulation of excessive plant nutrients, called eutrophication has occurred in Lake Erie, lake, lake Washington at Seattle, the lakes in Madison, area, Lake Zurich in Switzerland. An exhacutive study of urban water supply system in developing countries revealed that at least 60% of pollution is still dependent on underground source of drinking water, especially in other city or and distant villages, this very important source of water is now polluted from seepage bits, refuse dumps, septic tanks, barny and manures, transport accident and with diverse agricultural, chemical or biological pollutants. Otherwise important source of ground water pollution includes sewage and relable salts. The wide spread practice of dumping row sewage in shallow shark bit has made apprehansion of pollution of ground water in many cities. It altibutes in the rise of cholera, hepatitis, dysentery and other water borne disease, to the pollution of ground water especially in the zones where the water level is high. The most common complaint is susceptibility to worton- infection.

Oxygen deficiency also leads to production of objectionable odours in water. Due to the evaluation of putrefied gases, the solid wastes are buoyed up by these gases resulting in offensive foul matters floating on water surface. Depletion of oxygen content of water is caused by biological oxidation by microbes in any unit volume of water. BOD value is proportional to the amount of organic waste (oxidis able organic matter) present in water. Hence it is used as a measure of the degree of water pollution and waste level. Since the population of *E.coli* (Bacterium) increases tremendously, these start consuming most of the dissolved oxygen. Hence the dissolved oxygen content of water further decrease. The number of *E.coli* in the unit volume of water is also an

indicator for the degree of water pollution.

Accumulation of sewage and domestic wastes in water bodies retards the self regulatory capabilities of aquatic organisms. Self purifying ability of water is lost and it became unfit for domestic purposes. Moreover, the decomposition of wastes by aerobic microbes decreases due to the extreme pollution. In a balanced water ecosystem (Baig *et al.*, 2009; Mian *et al.*, 2010; Wang *et al.*, 2010), algae are found in small numbers. Most of these are beneficial because they enrich the oxygen content of water through the process the photosynthesis and are used as food also. When controlled quantity of domestic sewage is dumped into water, it becomes rich in nutrients (Bu *et al.*, 2010; Carpenter *et al.*, 1998) especially phosphates and nitrate. Thus, the water bodies become highly productive, i.e., eutrophic and the phenomenon is called eutrophication. A large portion of nutrients come back into water either when still alive or after their death or decay. As a result of this, the algae flora is shifted to blue-green algae, whose blooms are produced and can be seen floating as scums or blankets on the top water. These blooms forming algae are not utilized as food either by zooplankton or by invertebrates. Hence, they become pollutants themselves, because they are left to decompose and release their organic and inorganic matter back into water. The algae blooms compete with other aquatic plants for light to perform the process of photosynthesis. Thus oxygen level in depleted and CO₂ level gets increased. As result, fish and other animals begin to die and the clean water body is turned into a sinking drain. Thus eutrophication is a limiting factor in the supply of clean drinking water, for fishing and navigation etc.

¹Corresponding author

This paper reports physico-chemical characteristic, trace metal (Paul and Meyer, 2001; Akio, 1992; Friberg *et al.*, 1974) levels and phytoplankton composition in the reservoir. This is important because the reservoir serves as an alternative source of water supply to the municipal area in the dry season months (Jan-July) when there is water shortage.

MATERIALS AND METHODS

The important rural segments located between Jaunpur and Varanasi district named as Phoolpur Municipality has been under taken for investigations. Three sampling sites were chosen: site A is a drain that received waste water from market and residential area and some time sewage from a damaged septic tank nearby. Site B is at outflow of water containing animal wastes from dairy farm while Site C is the source of chemical effluents and is located behind the laboratories of college. At this site, the water is usually reddish brown and sandy substratum has the same colouration.

Water samples were collected fortnightly in acid washed polythene containers in 2005. Temperature was determined with 0-50°C thermometer while pH (Adetunde and Glover, 2010) and conductivity were determined using a portable conductivity meter and WPA conductivity meter. D.O. was determined by Winkler’s titrimetric method while ammonia-nitrogen and nitrate-nitrogen (Larry,

2006; Steiner *et al.*, 2006) were analysed according to usual methods (Bansal, 2018; Murali and Goel, 2014) (Atomic absorption spectrophotometer). Phytoplankton (John, 1989) were collected with a 64µm bolting silk plankton net and preserved in 4% formaline.

RESULTS

Table 1 shows the mean and ranges of physico-chemical parameters. D.O., pH, conductivity and ammonia-nitrogen showed wide variations during the study. The oxygen content for the reservoir is inadequate (mean 7.34±2.27 mgl⁻¹) and deoxygenation of the water occurred periodically. The pH is neutral (mean 7.27±0.32) while conductivity recorded intermediate value (mean 355.46±88.34 Scm⁻¹). Ammonia-nitrogen values were higher than nitrate-nitrogen values (Table-1). Iron is the most abundant trace metal in water (2.75±0.33 mgl⁻¹). Table-2 gives the composition and abundance of phytoplankton in the study area during the investigated period. The value obtained for Zn, Cu and Fe in the present study have been compared with WHO standards in Table-3. Generally, the blue-greens (Cynophyceae) dominated the community. There was a gradual increase in the abundance of Cynophytes from site A to C (Fig.1). Contrastingly, Bacillariophyceae and Desmidaceae decreased from site A to C. The green algae (Chlorophyceae) were second in order. Their abundance was relatively stable across site.

Table 1: Physico-chemical parameters in Phoolpur Municipality (Mean values±SD, range in parenthesis) in 2017


Parameters	Mean and Ranges
Temperature(°C)	34.2±2.35(16.2-52.4)
Dissolved Oxygen(mgl ⁻¹)	7.34±2.27(0.53-18.62)
pH	7.27±0.32(5.2-9.7)
Conductivity(µScm ⁻¹)	355.46±88.34(59.32-1088.82)
Ammonia-nitrogen(mgl ⁻¹)	0.93±0.37(0.01-2.87)
Nitrate-nitrogen(mgl ⁻¹)	0.75±0.08(0.01-0.47)
Zinc(mgl ⁻¹)	0.75±0.08(0.08-0.97)
Copper(mgl ⁻¹)	0.67±0.08(0.07-0.84)
Iron(mgl ⁻¹)	2.75±0.33(0.74-3.48)

Table 2: Phytoplankt on composition and abundance in Phoolpur Municipality

Phytoplankton	Total Abundance per cm ⁻³	% Abundance
Cynophyceae	450000	88.4
Chlorophyceae	63000	14.6
Desmidaceae	40000	8.2
Bacillariophyceae	15000	3.2
Dinophyceae	2000	0.4

Table 3: Mean trace metal levels in Phoolpur Municipal Area compared with WHO standards

Phytoplankton	Total Abundance per cm ⁻³	%Abundance
Zinc	0.34	6.00
Copper	0.48	2.00
Iron	2.37	0.70

**Figure 1: Phoolpur Municipal Area showing sampling sites**

DISCUSSION

The wide variations recorded in dissolved oxygen content, pH, conductivity and ammonia-nitrogen may be due to the fact that sampling took place during the late dry/rainy season periods. In tropical systems, marked variations in temperature and rainfall between seasons influence the physico- chemical characteristics of water bodies (Murali, 1995; Kaushal *et al.*, 2012). The conductivity of catchment area (Rida, 2017) can be recorded as being intermediate (Kennedy *et al.*, 1998). The lower nitrate-nitrogen values recorded relative to ammonia-nitrogen values has significant implications for nutrient. In addition, the Principal Component Analysis (PCA) identified as trace metals are the most important factor responsible for variations observed in the data. Although the levels of Zn and Cu are still below the recommended limits, there is need for urgent action to prevent further increase.

This is because trace metals act as micronutrients at low concentrations but at higher levels they become toxic depending on the prevalent chemical form (i.e., speciation) of trace element in water. Also bioaccumulation and food chain magnification of trace metals may occur (Mishra *et al.*, 2016).

However, the ferruginous nature of catchment area soils (Sharma *et al.*, 2017) may account for the levels of iron which exceeded recommended limits for drinking water quality. Cyanobacteria have higher requirements for

trace elements compared with eukaryotic phytoplankton (Yang, 2016; Valtcho, 2002; Aroroa *et al.*, 2002; Balaji *et al.*, 2022), hence increased level favours their growth and physiology.

Although phytoplankton species composition and diversity changes with environmental conditions such as nutrients levels, temperature, light, predator pressure etc., the relative importance of these factors vary considerably among different taxa. Under conditions of nutrient enrichment or eutrophication, the blue-greens are known to proliferate and form noxious blooms in fresh water environments (Hassan, 2013; Wang, 2020; Ehsanul Kabir *et al.*, 2012).

Cyano bacteria can produce both hepato toxic peptides which cause liver damage and are tumour-favouring e.g., microcystin as well as neurotoxic alkaloids that affect the nervous system e.g., aphonotoxin and anatoxin-a. Cyanophytes are thus recognized as potential toxic organisms and their blooms in drinking water are recreational reservoirs of great importance.

Cyano bacterial toxins are contained within the living cells and are not released into surrounding water until senescence or death of the cells. Pre-chlorination during water treatment or the application of copper sulphate in controlling blooms (used in some European Countries) results in cells death and release of toxins into water (Raluca *et al.*, 2017; Lakshmi *et al.*, 2022). In 1988 within 2-3 weeks of Cu treatment of the Goetzalkowice

reservoir, South Poland, the District Medical Officers recorded a rise in the number of cases of spastic bronchitis especially in children of this locality (Kumar *et al.*, 2005). Although there are no recorded cases of toxicosis in animals or humans (Prince *et al.*, 2020). Some abnormalities in the gonads of *Telapia zillii* and *Oreochromis niloticus* have been found in the reservoir. In *T. zillii*, abnormal ovaries were unusually shaped with very thin oocytes. However, in *O. niloticus* unequal ovarian lobes and the occurrence of two pairs of ovaries were observed. It was suggested that massive development of eggs could be due to break-down in nervous and hormonal co-ordination in fish during the breeding period. Since both species are algae feeders it is possible that the abnormalities observed could be related to their diet (Lal, 1996).

In view of the use of reservoir for water supply, control measures aimed at preventing bloom formation such as reduction of nutrient load into the reservoir, artificial mixing of the water and bio manipulation of food-web components should be implemented in order to protect bio-diversity (Feng *et al.*, 2008; Wang *et al.*, 2009) in India.

REFERENCES

- Akio M., 1992. Bitter Sea: The Human Cost of Minamata Disease. 1st Edition. Kosei Publishing, Japan.
- Adetunde L.A. and Glover R.L.K., 2010. Bacteriological Quality of Borehole Water Used by Students' of University for Development Studies, Navrongo Campus in Upper-East Region of Ghana. Current Research Journal of Biological Sciences, **2**(6): 361-364.
- Arora A., Pabby A. and Singh P.K., 2002. Accumulation of heavy metals by *Azolla* species. The annual conferences on contaminated soils and waters. University of Massachusetts, U.S.A.
- Balaji L., Muthukannan M. and Kanniga Dev R., 2022. A GIS-Based Study of Air and Water Quality Trends in Madurai City, India. Nature Environment and Pollution Technology An International Quarterly Scientific Journal, **21**(1): 21-32.
- Bansal N., 2018. Industrial Development and Challenges of Water Pollution in Coastal Areas: The Case of Surat, India IOP Conf. Series: Earth and Environmental Science 120.
- Baig J.A., Kazi T.G., Arain M.B., Afridi H.I., Kandhro G.A., Sarfraz R.A., Jamali M.K. and Shah A.Q., 2009. Evaluation of arsenic and other physico-chemical parameters of surface and ground water of Jamshoro, Pakistan. Journal of Hazardous Materials, **166**: 662–669.
- Bu H., Tan X., Li S. and Zhang Q., 2010. Water quality assessment of the Jinshui River (China) using multivariate statistical techniques. Environ Earth Sci., **60**: 1631–1639.
- Carpenter S.R., Caraco N.F., Correll D.L., Howarth R.W., Sharpley A.N. and Smith V.H., 1998. Non point pollution of surface waters with phosphorus and nitrogen. Ecological Applications, **8**: 559-568.
- Feng M.Q., Zheng B.M. and Zhou X.D., 2008. Water Environment Simulation and Prediction, Science Press, Beijing, China.
- Friberg L., Piscator M., Nordberg G.F and Kjellstrom T., 1974. Cadmium in the environment, 2nd edition, Chemical Rubber Company Press, Cleveland, Ohio, pp. 248.
- Goudar C.T., Ganji S.H., Pujar B.G. and Strevett K.A., 2000. Substrate inhibition Kinetics of Phenol Biodegradation, J.WEF WEF., **72**(1): 50-55.
- Hassan R., Hassan A., Montasser M. and Latif N.A., 2013. Synthesis of New Schiff Base from Natural Products for Remediation of Water Pollution with Heavy Metals in Industrial Areas” Journal of Chemistry.
- John H. Seinfeld, 1989. Urban Air Pollution: State of the Science. Science, **243**(4892): 745-752.
- Kaushal R.K., George K. Varghese and Chabukdhara M., 2012. Studied on Municipal Solid Waste Management in India-Current State and Future Challenges: A Review. Rajendra Kumar Kaushal et al. / International Journal of Engineering Science and Technology (IJEST), **4**(4).
- Kabir E., Ray S., Kim K.-H., Yoon H.-O., Jeon E.-C., Kim Y.S., Cho Y.-S., Yun S.-T. and Brown R.J.C., 2012. “Current Status of Trace Metal Pollution in Soils Affected by Industrial Activities”, The Scientific World Journal.
- Khan A.M., 1998. Studies & Evaluation of Physico-Chemical parameters in Sai River, Ph.D. Thesis.
- Kennedy V.J., Augusthy A., Varier K.M., Magudapathy P., Panchapakesan S., Ramesh C., Nair K.G.M. and Vijayan V. 1998. PIXE analysis of trace pollutants in Chaliyar river water in Malabar, India. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions

- with Materials and Atoms, **134**(2): 224-228.
- Lal A.K., 1996. Environmental status of Delhi. Indian Journal of Environmental Protection, **16**(1): 1–11.
- Lakshmi T. and Veeraragavan G.R., 2022. Conducted a research on “A Survey on Nanotechnology-Based Bioremediation of Wastewater” Journal of Bioinorganic Chemistry and Applications.
- Larry W., 2006. World Water Day. A Billion people Worldwide Lack Safe Drinking <http://environment.about.com/od/environmentalevents/a/waterdayqa.htm>, March 22nd, 2006.
- Mian I.A., Begum S., Riaz M., Ridealgh M., Mc Clean C.J. and Cresser M.S., 2010. Spatial and temporal trends in nitrate concentrations in the River Derwent, North Yorkshire, and its need for NVZ status. Science of the Total Environment, **408**: 702–712.
- Murali Krishna Gurram and Geol Geosci J., 2014. An Appraisal of Sustainability Scenario of Solid Waste Management: AGIS Study on Municipal Wards of Hyderabad, India”, Journal of Geology & Geosciences, **3**: 2.
- Murali Krishna K.V.S.G., 1995. Air pollution and control, Kaushal & Co., Kakinada (A.P), India, pp. 307.
- Mishra S., Kumar A. and Shukla P., 2016. Study of water quality in Hindon River using pollution index and environmetrics, India. Desalination Water Treat, **57**: 19121-19130.
- Raluca M.H., Maria G., Teresa T., Lidia F. and Giuseppe O., 2017. Studied on “Bioremediation: An Overview on Current Practices, Advances, and New Perspectives in Environmental Pollution Treatment”. Bio. Med. Research International.
- Rida Al-Adamat, 2017. Modelling Surface Water Susceptibility to Pollution Using GIS. Journal of Geographic Information System, **9**(3): 293-308.